arXiv:1008.4048v1 [math.NT] 24 Aug 2010

ON A CONJECTURE CONCERNING THE FRACTION OF INVERTIBLE M-TIMES PERSYMMETRIC MATRICES OVER $\mathbb{F}_2$

JORGEN CHERLY

Résumé. Dans cette article nous annonçons que la fraction d'invertible m-fois matrices persymétriques sur $\mathbb{F}_2$ est égale á $\prod_{j=1}^{m}(1-2^{-j})$

Abstract. In this paper we state that the fraction of invertible m-times persymmetric matrices over $\mathbb{F}_2$ is equal to $\prod_{j=1}^{m}(1-2^{-j})$

Contents

1. Some notations concerning m-times persymmetric matrices over $\mathbb{F}_2$ 3
2. A conjecture concerning m-times persymmetric matrices 7
3. Justification of our assumption (2.1) in the first case 7
4. Justification of our assumption (2.1) in the second case 7
5. Justification of our assumption (2.1) in the third case 8
6. Justification of our assumption (2.1) in the fourth case 8
7. Justification of our assumption (2.1) in the fifth case 8
8. An example 9

References 11

1. Some notations concerning m-times persymmetric matrices over $\mathbb{F}_2$

Let $m, s_1, s_2, \ldots s_m, \delta, k$ be positive integers such that $1 \leqslant m \leqslant \sum_{j=1}^{m} s_j = \delta \leqslant k$.
We can assume without restriction that $1 \leqslant s_1 \leqslant s_2 \leqslant \ldots \leqslant s_m$ since the rank of a matrix does not change under elementary row operations.
We denote by $\Gamma_{\delta}^{\left[\begin{smallmatrix} s_1 \\ \vdots \\ s_m \end{smallmatrix}\right] \times k}$ the number of rank δ m- times persymmetric matrices over the finite field $\mathbb{F}_2$ of the below form

$$
\left(\begin{array}{ccccc}
\alpha_1^{(1)} & \alpha_2^{(1)} & \ldots & \alpha_{k-1}^{(1)} & \alpha_k^{(1)} \\
\alpha_2^{(1)} & \alpha_3^{(1)} & \ldots & \alpha_k^{(1)} & \alpha_{k+1}^{(1)} \\
\vdots & \vdots & \vdots & \vdots & \vdots \\
\alpha_{s_1}^{(1)} & \alpha_{s_1+1}^{(1)} & \ldots & \alpha_{s_1+k-2}^{(1)} & \alpha_{s_1+k-1}^{(1)} \\
\hline
\alpha_1^{(2)} & \alpha_2^{(2)} & \ldots & \alpha_{k-1}^{(2)} & \alpha_k^{(2)} \\
\alpha_2^{(2)} & \alpha_3^{(2)} & \ldots & \alpha_k^{(2)} & \alpha_{k+1}^{(2)} \\
\vdots & \vdots & \vdots & \vdots & \vdots \\
\alpha_{s_2}^{(2)} & \alpha_{s_2+1}^{(2)} & \ldots & \alpha_{s_2+k-2}^{(2)} & \alpha_{s_2+k-1}^{(2)} \\
\hline
\alpha_1^{(3)} & \alpha_2^{(3)} & \ldots & \alpha_{k-1}^{(3)} & \alpha_k^{(3)} \\
\alpha_2^{(3)} & \alpha_3^{(3)} & \ldots & \alpha_k^{(3)} & \alpha_{k+1}^{(3)} \\
\vdots & \vdots & \vdots & \vdots & \vdots \\
\alpha_{s_3}^{(3)} & \alpha_{s_3+1}^{(3)} & \ldots & \alpha_{s_3+k-2}^{(3)} & \alpha_{s_3+k-1}^{(3)} \\
\hline
\vdots & \vdots & \vdots & \vdots & \vdots \\
\hline
\alpha_1^{(m)} & \alpha_2^{(m)} & \ldots & \alpha_{k-1}^{(m)} & \alpha_k^{(m)} \\
\alpha_2^{(m)} & \alpha_3^{(m)} & \ldots & \alpha_k^{(m)} & \alpha_{k+1}^{(m)} \\
\vdots & \vdots & \vdots & \vdots & \vdots \\
\alpha_{s_m}^{(m)} & \alpha_{s_m+1}^{(m)} & \ldots & \alpha_{s_m+k-2}^{(m)} & \alpha_{s_m+k-1}^{(m)}
\end{array}\right) \tag{1.1}
$$

$\boxed{\textbf{The case } s_1 = s_2 = \ldots = s_m = 1,\ m = \delta \leqslant k}$

Then $\delta = m$ and $\Gamma_{\delta}^{\left[\begin{smallmatrix} 1 \\ \vdots \\ 1 \end{smallmatrix}\right] \times k}$ denotes the number of rank m $m \times k$ matrices over $\mathbb{F}_2$ of the below form

$$\begin{pmatrix}
\alpha_1^{(1)} & \alpha_2^{(1)} & \ldots & \alpha_{k-1}^{(1)} & \alpha_k^{(1)} \\
\alpha_1^{(2)} & \alpha_2^{(2)} & \ldots & \alpha_{k-1}^{(2)} & \alpha_k^{(2)} \\
\alpha_1^{(3)} & \alpha_2^{(3)} & \ldots & \alpha_{k-1}^{(3)} & \alpha_k^{(3)} \\
\vdots & \vdots & \vdots & \vdots & \vdots \\
\alpha_1^{(m)} & \alpha_2^{(m)} & \ldots & \alpha_{k-1}^{(m)} & \alpha_k^{(m)}
\end{pmatrix} \tag{1.2}$$

$\boxed{\textbf{The case } m = 1, \quad 2 \leqslant s_1 = \delta \leqslant k}$

Then $\delta = s_1$ and $\Gamma_\delta^{[s_1]\times k}$denotes the number of persymmetric $s_1 \times k$ rank s_1 matrices over $\mathbb{F}_2$ of the below form

$$\begin{pmatrix}
\alpha_1^{(1)} & \alpha_2^{(1)} & \ldots & \alpha_{k-1}^{(1)} & \alpha_k^{(1)} \\
\alpha_2^{(1)} & \alpha_3^{(1)} & \ldots & \alpha_k^{(1)} & \alpha_{k+1}^{(1)} \\
\vdots & \vdots & \vdots & \vdots & \vdots \\
\alpha_{s_1}^{(1)} & \alpha_{s_1+1}^{(1)} & \ldots & \alpha_{s_1+k-2}^{(1)} & \alpha_{s_1+k-1}^{(1)}
\end{pmatrix} \tag{1.3}$$

$\boxed{\textbf{The case } m = 2, \quad 2 \leqslant s_1 \leqslant s_2, \ s_1 + s_2 = \delta \leqslant k}$

Then $\delta = s_1 + s_2$ and $\Gamma_\delta^{\left[\substack{s_1 \\ s_2}\right]\times k}$denotes the number of double persymmetric

$(s_1 + s_2) \times k$ rank $s_1 + s_2$ matrices over $\mathbb{F}_2$ of the below form

$$\left(\begin{array}{ccccc}
\alpha_1^{(1)} & \alpha_2^{(1)} & \ldots & \alpha_{k-1}^{(1)} & \alpha_k^{(1)} \\
\alpha_2^{(1)} & \alpha_3^{(1)} & \ldots & \alpha_k^{(1)} & \alpha_{k+1}^{(1)} \\
\vdots & \vdots & \vdots & \vdots & \vdots \\
\alpha_{s_1}^{(1)} & \alpha_{s_1+1}^{(1)} & \ldots & \alpha_{s_1+k-2}^{(1)} & \alpha_{s_1+k-1}^{(1)} \\
\hline
\alpha_1^{(2)} & \alpha_2^{(2)} & \ldots & \alpha_{k-1}^{(2)} & \alpha_k^{(2)} \\
\alpha_2^{(2)} & \alpha_3^{(2)} & \ldots & \alpha_k^{(2)} & \alpha_{k+1}^{(2)} \\
\vdots & \vdots & \vdots & \vdots & \vdots \\
\alpha_{s_2}^{(2)} & \alpha_{s_2+1}^{(2)} & \ldots & \alpha_{s_2+k-2}^{(2)} & \alpha_{s_2+k-1}^{(2)}
\end{array}\right) \tag{1.4}$$

$\boxed{\textbf{The case } m = 3, \quad 2 \leqslant s_1 \leqslant s_2 \leqslant s_3, \ s_1 + s_2 + s_3 = \delta \leqslant k}$

Then $\delta = s_1 + s_2 + s_3$ and $\Gamma_\delta^{\left[\substack{s_1 \\ s_2 \\ s_3}\right]\times k}$ denotes the number of triple persymmetric
$(s_1 + s_2 + s_3) \times k$ rank $s_1 + s_2 + s_3$ matrices over $\mathbb{F}_2$ of the below form

$$
\left(\begin{array}{ccccc}
\alpha_1^{(1)} & \alpha_2^{(1)} & \ldots & \alpha_{k-1}^{(1)} & \alpha_k^{(1)} \\
\alpha_2^{(1)} & \alpha_3^{(1)} & \ldots & \alpha_k^{(1)} & \alpha_{k+1}^{(1)} \\
\vdots & \vdots & \vdots & \vdots & \vdots \\
\alpha_{s_1}^{(1)} & \alpha_{s_1+1}^{(1)} & \ldots & \alpha_{s_1+k-2}^{(1)} & \alpha_{s_1+k-1}^{(1)} \\
\hline
\alpha_1^{(2)} & \alpha_2^{(2)} & \ldots & \alpha_{k-1}^{(2)} & \alpha_k^{(2)} \\
\alpha_2^{(2)} & \alpha_3^{(2)} & \ldots & \alpha_k^{(2)} & \alpha_{k+1}^{(2)} \\
\vdots & \vdots & \vdots & \vdots & \vdots \\
\alpha_{s_2}^{(2)} & \alpha_{s_2+1}^{(2)} & \ldots & \alpha_{s_2+k-2}^{(2)} & \alpha_{s_2+k-1}^{(2)} \\
\hline
\alpha_1^{(3)} & \alpha_2^{(3)} & \ldots & \alpha_{k-1}^{(3)} & \alpha_k^{(3)} \\
\alpha_2^{(3)} & \alpha_3^{(3)} & \ldots & \alpha_k^{(3)} & \alpha_{k+1}^{(3)} \\
\vdots & \vdots & \vdots & \vdots & \vdots \\
\alpha_{s_3}^{(3)} & \alpha_{s_3+1}^{(3)} & \ldots & \alpha_{s_3+k-2}^{(3)} & \alpha_{s_3+k-1}^{(3)}
\end{array}\right) \tag{1.5}
$$

$$\boxed{\textbf{The case } m \geqslant 4, \quad s_1 = s_2 = \ldots = s_{m-3} = 1,\ 2 \leqslant s_{m-2} \leqslant s_{m-1} \leqslant s_m,\ \sum_{j=1}^{m} s_j = \delta \leqslant k}$$

Then $\delta = m-3+s_{m-2}+s_{m-1}+s_m$ and $\Gamma_{\delta}^{\left[\begin{smallmatrix}1\\ \vdots\\ 1\\ s_{m-2}\\ s_{m-1}\\ s_m\end{smallmatrix}\right]\times k}$ denotes the number of $m-times$ persymmetric $\delta \times k$ rank δ matrices over $\mathbb{F}_2$ of the below form

$$
\left(\begin{array}{ccccc}
\alpha_1^{(1)} & \alpha_2^{(1)} & \cdots & \alpha_{k-1}^{(1)} & \alpha_k^{(1)} \\
\alpha_1^{(2)} & \alpha_2^{(2)} & \cdots & \alpha_{k-1}^{(2)} & \alpha_k^{(2)} \\
\alpha_1^{(3)} & \alpha_2^{(3)} & \cdots & \alpha_{k-1}^{(3)} & \alpha_k^{(3)} \\
\vdots & \vdots & \vdots & \vdots & \vdots \\
\alpha_1^{(m-3)} & \alpha_2^{(m-3)} & \cdots & \alpha_{k-1}^{(m-3)} & \alpha_k^{(m-3)} \\
\hline
\alpha_1^{(m-2)} & \alpha_2^{(m-2)} & \cdots & \alpha_{k-1}^{(m-2)} & \alpha_k^{(m-2)} \\
\alpha_2^{(m-2)} & \alpha_3^{(m-2)} & \cdots & \alpha_k^{(m-2)} & \alpha_{k+1}^{(m-2)} \\
\vdots & \vdots & \vdots & \vdots & \vdots \\
\alpha_{s_{m-2}}^{(m-2)} & \alpha_{s_{m-2}+1}^{(m-2)} & \cdots & \alpha_{s_{m-2}+k-2}^{(m-2)} & \alpha_{s_{m-2}+k-1}^{(m-2)} \\
\hline
\alpha_1^{(m-1)} & \alpha_2^{(m-1)} & \cdots & \alpha_{k-1}^{(m-1)} & \alpha_k^{(m-1)} \\
\alpha_2^{(m-1)} & \alpha_3^{(m-1)} & \cdots & \alpha_k^{(m-1)} & \alpha_{k+1}^{(m-1)} \\
\vdots & \vdots & \vdots & \vdots & \vdots \\
\alpha_{s_{m-1}}^{(m-1)} & \alpha_{s_{m-1}+1}^{(m-1)} & \cdots & \alpha_{s_{m-1}+k-2}^{(m-1)} & \alpha_{s_{m-1}+k-1}^{(m-1)} \\
\hline
\alpha_1^{(m)} & \alpha_2^{(m)} & \cdots & \alpha_{k-1}^{(m)} & \alpha_k^{(m)} \\
\alpha_2^{(m)} & \alpha_3^{(m)} & \cdots & \alpha_k^{(m)} & \alpha_{k+1}^{(m)} \\
\vdots & \vdots & \vdots & \vdots & \vdots \\
\alpha_{s_m}^{(m)} & \alpha_{s_m+1}^{(m)} & \cdots & \alpha_{s_m+k-2}^{(m)} & \alpha_{s_m+k-1}^{(m)}
\end{array}\right) \tag{1.6}
$$

2. A CONJECTURE CONCERNING M-TIMES PERSYMMETRIC MATRICES

Conjecture : Let $1 \leqslant m \leqslant \sum_{j=1}^{m} s_j = \delta \leqslant k$.

We state that the number $\Gamma_{\delta}^{\left[\begin{smallmatrix} s_1 \\ \vdots \\ s_m \end{smallmatrix}\right]\times k}$ of rank δ m-times persymmetric matrices over the finite field $\mathbb{F}_2$ of the form (1.1) is equal to

$$2^{\delta-m}\prod_{j=1}^{m}(2^k - 2^{\delta-j}) = 2^{(1+m)\delta-\frac{m^2}{2}-\frac{3m}{2}}\prod_{j=1}^{m}(2^{k-\delta+j}-1) \tag{2.1}$$

In the case $\delta = k$ we deduce easily from (2.1) that the fraction of invertible m-times persymmetric matrices is equal to $\prod_{j=1}^{m}(1-2^{-j})$.
To justify our assumption (2.1) we shall prove that our supposition is valid in the following five cases:

- (1) **The case** $s_1 = s_2 = \ldots = s_m = 1,\ m = \delta \leqslant k$
- (2) **The case** $m = 1, \quad 2 \leqslant s_1 = \delta \leqslant k$
- (3) **The case** $m = 2, \quad 2 \leqslant s_1 \leqslant s_2,\ s_1 + s_2 = \delta \leqslant k$
- (4) **The case** $m = 3, \quad 2 \leqslant s_1 \leqslant s_2 \leqslant s_3,\ s_1 + s_2 + s_3 = \delta \leqslant k$
- (5) **The case** $m \geqslant 4, \quad s_1 = s_2 = \ldots = s_{m-3} = 1,\ 2 \leqslant s_{m-2} \leqslant s_{m-1} \leqslant s_m,\ \sum_{j=1}^{m} s_j = \delta \leqslant k$

3. JUSTIFICATION OF OUR ASSUMPTION (2.1) IN THE FIRST CASE

Theorem 3.1. *The number of* $m \times k$ *matrices over* $\mathbb{F}_2$ *of rank* m *is equal to* $\prod_{l=0}^{m-1}(2^k - 2^l) = \prod_{j=1}^{m}(2^k - 2^{m-j})$

Proof. See (1.2) refer to Section 1.

We have, $\Gamma_{m}^{\left[\begin{smallmatrix} 1 \\ \vdots \\ 1 \end{smallmatrix}\right]\times k} = \prod_{l=0}^{m-1}(2^k - 2^l)$ (See George Landsberg [1] or S.D. Fisher and M.N Alexander [2]) ☐

4. JUSTIFICATION OF OUR ASSUMPTION (2.1) IN THE SECOND CASE

Theorem 4.1. *The number of* $\delta \times k$ *persymmetric matrices over* $\mathbb{F}_2$ *of rank* $\delta = s_1$ *is equal to* $2^{k+\delta-1} - 2^{2\delta-2} = 2^{\delta-1}\prod_{j=1}^{1}(2^k - 2^{\delta-j})$

Proof. See (1.3) refer to Section 1.
We have,
$\Gamma_{\delta}^{[s_1]\times k} = 2^{k+\delta-1} - 2^{2\delta-2}$ (See David E. Daykin [3] or Cherly [4]) $\square$

5. Justification of our assumption (2.1) in the third case

Theorem 5.1. *The number of $\delta \times k$ double persymmetric matrices over $\mathbb{F}_2$ of rank $\delta = s_1 + s_2$ is equal to* $2^{\delta-2}\prod_{j=1}^{2}(2^k - 2^{\delta-j})$

Proof. See (1.4) refer to Section 1.
We have,
$\Gamma_{\delta}^{\left[\begin{smallmatrix} s_1 \\ s_2 \end{smallmatrix}\right]\times k} = 2^{2k+\delta-2} - 3\cdot 2^{k+2\delta-4} + 2^{3\delta-5} = 2^{\delta-2}(2^{2k} - 3\cdot 2^{k+\delta-2} + 2^{2\delta-3}) = 2^{\delta-2}\prod_{j=1}^{2}(2^k - 2^{\delta-j})$ (See Cherly [5]) $\square$

6. Justification of our assumption (2.1) in the fourth case

Theorem 6.1. *The number of $\delta \times k$ triple persymmetric matrices over $\mathbb{F}_2$ of rank $\delta = s_1 + s_2 + s_3$ is equal to* $2^{\delta-3}\prod_{j=1}^{3}(2^k - 2^{\delta-j})$

Proof. See (1.5) refer to Section 1.
We have,
$\Gamma_{\delta}^{\left[\begin{smallmatrix} s_1 \\ s_2 \\ s_3 \end{smallmatrix}\right]\times k} = 2^{3k+\delta-3} - 7\cdot 2^{2k+2\delta-6} + 7\cdot 2^{k+3\delta-8} + 2^{4\delta-9} = 2^{\delta-3}(2^{3k} - 7\cdot 2^{2k+2\delta-3} + 7\cdot 2^{k+2\delta-5} - 2^{3\delta-6}) = 2^{\delta-3}\prod_{j=1}^{3}(2^k - 2^{\delta-j})$ (See Cherly [6]) $\square$

7. Justification of our assumption (2.1) in the fifth case

Theorem 7.1. *The number of $\delta \times k$ m-times persymmetric matrices over $\mathbb{F}_2$ of rank $\delta = m - 3 + s_1 + s_2 + s_3$ is equal to* $2^{\delta-m}\prod_{j=1}^{m}(2^k - 2^{\delta-j})$

Proof. See (1.6) refer to Section 1.
We obtain from Lemma 10.1 in [4] with some obviously modifications that

$$\Gamma_{\delta}^{\left[\begin{smallmatrix} 1 \\ \vdots \\ 1 \\ s_{m-2} \\ s_{m-1} \\ s_m \end{smallmatrix}\right]\times k} = \Gamma_{\delta-m+3}^{\left[\begin{smallmatrix} s_{m-2} \\ s_{m-1} \\ s_m \end{smallmatrix}\right]\times k}\prod_{j=1}^{m-3}(2^k - 2^{\delta-j}) \tag{7.1}$$

We get from Theorem 6.1 that

$$(7.2)\quad \Gamma_{\delta-m+3}^{\left[\begin{smallmatrix} s_{m-2}\\ s_{m-1}\\ s_m\end{smallmatrix}\right]\times k} = 2^{\delta-m+3-3}\prod_{j=1}^{3}(2^k-2^{\delta-m+3-j}) = 2^{\delta-m}\prod_{j=m-2}^{m}(2^k-2^{\delta-j})$$

Combining (7.1) and (7.2) we obtain

$$\Gamma_{\delta}^{\left[\begin{smallmatrix} 1\\ \vdots\\ 1\\ s_{m-2}\\ s_{m-1}\\ s_m\end{smallmatrix}\right]\times k} = 2^{\delta-m}\prod_{j=1}^{m}(2^k-2^{\delta-j}) \qquad \square$$

For other related articles concerning persymmetric matrices over the finite field with two elements see Cherly [7],[8] and [9].

8. An example

Let m, k, δ be rational integers such that $1 \leqslant m \leqslant \delta \leqslant k$.
Let $\delta \equiv i (mod\, m)$ that is $\delta = i + s \cdot m$ such that $0 \leqslant i \leqslant m-1$.

We denote by $M_{m,\delta,k}$ the $k \times \delta$ matrix over $\mathbb{F}_2$ of the below form:

$$(8.1)\quad \left(\begin{array}{ccccc|cccc}
\alpha_1 & \alpha_2 & \alpha_3 & \cdots & \alpha_m & \alpha_{m+1} & \alpha_{m+2} & \cdots & \alpha_\delta \\
\alpha_{m+1} & \alpha_{m+2} & \alpha_{m+3} & \cdots & \alpha_{2m} & \alpha_{2m+1} & \alpha_{2m+2} & \cdots & \alpha_{\delta+m} \\
\alpha_{2m+1} & \alpha_{2m+2} & \alpha_{2m+3} & \cdots & \alpha_{3m} & \alpha_{3m+1} & \alpha_{3m+2} & \cdots & \alpha_{\delta+2m} \\
\vdots & \vdots & \vdots & \vdots & \vdots & \vdots & \vdots & \vdots & \vdots \\
\vdots & \vdots & \vdots & \vdots & \vdots & \vdots & \vdots & \vdots & \vdots \\
\alpha_{(m-1)m+1} & \alpha_{(m-1)m+2} & \alpha_{(m-1)m+3} & \cdots & \alpha_{m^2} & \alpha_{m^2+1} & \alpha_{m^2+2} & \cdots & \alpha_{\delta+(m-1)m} \\
\hline
\alpha_{m^2+1} & \alpha_{m^2+2} & \alpha_{m^2+3} & \cdots & \alpha_{m^2+m} & \alpha_{m^2+m+1} & \alpha_{m^2+m+2} & \cdots & \alpha_{\delta+m^2} \\
\alpha_{m^2+m+1} & \alpha_{m^2+m+2} & \alpha_{m^2+m+3} & \cdots & \alpha_{m^2+2m} & \alpha_{m^2+2m+1} & \alpha_{m^2+2m+2} & \cdots & \alpha_{\delta+m^2+m} \\
\vdots & \vdots & \vdots & \vdots & \vdots & \vdots & \vdots & \vdots & \vdots \\
\vdots & \vdots & \vdots & \vdots & \vdots & \vdots & \vdots & \vdots & \vdots \\
\alpha_{(k-1)m+1} & \alpha_{(k-1)m+2} & \alpha_{(k-1)m+3} & \cdots & \alpha_{km} & \alpha_{km+1} & \alpha_{km+2} & \cdots & \alpha_{\delta+(k-1)m}
\end{array}\right).$$

The transpose of $M_{m,\delta,k}$ becomes after a rearrangement of the rows a m-times persymmetric $\delta \times k$ matrix with $s_1 = s_2 = \ldots = s_{m-i} = s$ and $s_{m-i+1} = s_{m-i+2} = \ldots s_m = s+1$, refer to Section 1.
We then deduce from the conjecture (2.1) that the number of rank δ matrices of the form (8.1) is equal to $2^{\delta-m}\prod_{j=1}^{m}(2^k-2^{\delta-j})$

Example. Let m=3, $\delta = 8$ and k=10.
$8 \equiv 2 (mod\; 3)$ that is $8 = 2 + 2 \cdot 3$, so i=2 and s=2.

Then $M_{3,8,10}$ denotes the 10×8 matrix over $\mathbb{F}_2$ of the below form:

$$\left(\begin{array}{ccc|ccccc}
\alpha_1 & \alpha_2 & \alpha_3 & \alpha_4 & \alpha_5 & \alpha_6 & \alpha_7 & \alpha_8 \\
\alpha_4 & \alpha_5 & \alpha_6 & \alpha_7 & \alpha_8 & \alpha_9 & \alpha_{10} & \alpha_{11} \\
\alpha_7 & \alpha_8 & \alpha_9 & \alpha_{10} & \alpha_{11} & \alpha_{12} & \alpha_{13} & \alpha_{14} \\
\hline
\alpha_{10} & \alpha_{11} & \alpha_{12} & \alpha_{13} & \alpha_{14} & \alpha_{15} & \alpha_{16} & \alpha_{17} \\
\alpha_{13} & \alpha_{14} & \alpha_{15} & \alpha_{16} & \alpha_{17} & \alpha_{18} & \alpha_{19} & \alpha_{20} \\
\alpha_{16} & \alpha_{17} & \alpha_{18} & \alpha_{19} & \alpha_{20} & \alpha_{21} & \alpha_{22} & \alpha_{23} \\
\alpha_{19} & \alpha_{20} & \alpha_{21} & \alpha_{22} & \alpha_{23} & \alpha_{24} & \alpha_{25} & \alpha_{26} \\
\alpha_{22} & \alpha_{23} & \alpha_{24} & \alpha_{25} & \alpha_{26} & \alpha_{27} & \alpha_{28} & \alpha_{29} \\
\alpha_{25} & \alpha_{26} & \alpha_{27} & \alpha_{28} & \alpha_{29} & \alpha_{30} & \alpha_{31} & \alpha_{32} \\
\alpha_{28} & \alpha_{29} & \alpha_{30} & \alpha_{31} & \alpha_{32} & \alpha_{33} & \alpha_{34} & \alpha_{35}
\end{array}\right). \tag{8.2}$$

We then obtain that the number $\Gamma_8^{\left[\begin{smallmatrix}2\\3\\3\end{smallmatrix}\right]\times 10}$ of rank $\delta = 8$ matrices of the form (8.2) is equal to $2^5\prod_{j=1}^{3}(2^{10} - 2^{8-j}) = 2^{23} \cdot 7 \cdot 15 \cdot 31 = 3255 \cdot 2^{23}$.

References

[1] Landsberg, G Ueber eine Anzahlbestimmung und eine damit zusammenhangende Reihe, J. reine angew.Math, **111**(1893),87-88.

[2] Fisher,S.D and Alexander M.N. Matrices over a finite field
Amer.Math.Monthly 73(1966), 639-641

[3] Daykin David E, Distribution of Bordered Persymmetric Matrices in a finite field

[4] Cherly, Jorgen.
Exponential sums and rank of persymmetric matrices over $\mathbf{F}_2$
arXiv : 0711.1306, 46 pp

[5] Cherly, Jorgen.
Exponential sums and rank of double persymmetric matrices over $\mathbf{F}_2$
arXiv : 0711.1937, 160 pp

[6] Cherly, Jorgen.
Exponential sums and rank of triple persymmetric matrices over $\mathbf{F}_2$
arXiv : 0803.1398, 233 pp

[7] Cherly, Jorgen.
Results about persymmetric matrices over $\mathbf{F}_2$ and related exponentials sums
arXiv : 0803.2412v2, 32 pp

[8] Cherly, Jorgen.
Polynomial equations and rank of matrices over $\mathbf{F}_2$ related to persymmetric matrices
arXiv : 0909.0438v1, 33 pp

[9] Cherly, Jorgen.
On a conjecture regarding enumeration of n-times persymmetric matrices over $\mathbf{F}_2$ by rank
arXiv : 0909.4030, 21 pp

Département de Mathématiques, Université de Brest, 29238 Brest cedex 3, France
E-mail address: `Jorgen.Cherly@univ-brest.fr`
E-mail address: `andersen69@wanadoo.fr`